Chapter 12 - Liquids and Solids

12-1 Liquids

- I. Properties of Liquids and the Kinetic Molecular Theory
 - A. Fluids
 - 1. Substances that can flow and therefore take the shape of their container
 - B. Relative High Density
 - 1. 10% less dense than solids (average)
 - a. Water is an exception
 - 2. 1000x more dense than gases
 - C. Relative Incompressibility
 - 1. The volume of liquids doesn't change appreciably when pressure is applied
 - D. Ability to Diffuse
 - 1. Liquids diffuse and mix with other liquids
 - 2. Rate of diffusion increases with temperature (↑ average Kinetic Energy)
 - E. Surface Tension
 - 1. Surface Tension
 - a. A force that tends to pull adjacent parts of a liquid's surface together, thereby decreasing surface area to the smallest possible size
 - b. Hydrogen bonding in water creates stronger than normal surface tension
 - 2. Capillary Action
 - a. The attraction of the surface of a liquid to the surface of a solid
 - F. Evaporation and Boiling
 - 1. Vaporization
 - a. The process by which a liquid of solid changes to a gas
 - 2. Evaporation
 - a. The process by which particles escape from the surface of a nonboiling liquid enter the gas state
 - b. Evaporation is a form of vaporization
 - 3. Boiling
 - a. The change of a liquid to bubbles of vapor that appear throughout the liquid
 - G. Formation of Solids
 - 1. Freezing (or Solidification)
 - a. The physical change of a liquid to a solid by removal of heat

Solid Liquid Gas

12-2 Solids

- I. Properties of Solids and the Kinetic Molecular Theory
 - A. Types of Solids
 - 1. Crystalline Solids substances in which the particles are arranged in an orderly, geometric, repeating pattern
 - 2. Amorphous Solids substances in which the particles are arranged randomly
 - B. Definite Shape and Volume
 - C. Definite Melting Point
 - 1. Melting is the physical change of a solid to a liquid by the addition of heat
 - 2. Melting point is the temperature at which a solid becomes a liquid
 - a. Crystalline solids have definite melting points
 - b. Amorphous solids do not have definite melting points
 - D. High Density and Incompressibility
 - E. Low Rate of Diffusion
 - 1. Two solids in contact will experience VERY SLOW rates of diffusion
- II. Crystalline Solids
 - A. Crystal Structure
 - 1. The total three dimensional arrangement of particles of a crystal
 - B. Unit Cell
 - 1. The smallest portion of a crystal lattice that shows the three-dimensional pattern of the entire lattice







lsometric or cubic

Tetragonal

Hexagonal

т







Monoclinic

- C. Binding Forces in Solids
 - 1. Ionic crystals
 - 2. Covalent network crystals a. Diamond, guartz
 - 3. Metallic crystals
 - 4. Covalent Molecular crystals
 - a. Ice
- III. Amorphous Solids
 - A. "Amorphous"
 - 1. Greek for "without shape"
 - B. Formation of amorphous solids
 - 1. Rapid cooling of molten materials can prevent the formation of crystals
 - a. Glass
 - b. Obsidian

12-3 Changes of State

- <u>Equilibrium</u>
 - A. Equilibrium
 - 1. Dynamic condition in which two opposing changes occur at equal rates in a closed system
 - B. Equilibrium and Changes of State
 - 1. Phase
 - a. Any part of a system that has uniform composition and properties
 - 2. Condensation
 - a. The process by which a gas changes to a liquid
 - 3. A closed system at constant temperature will reach an equilibrium position at which the rates of evaporation and condensation will be the same
 - C. An Equilibrium Equation
 - liquid + heat enegy \leftrightarrow vapor
 - D. Le Chatelier's Principle

1.

- 1. When a system at equilibrium is disturbed by application of a stress, it attains a new equilibrium position that minimizes the stress
- E. Equilibrium and Temperature
 - 1. Increasing the temperature will move the more particles into the vapor phase to compensate for the new energy



25 Celsius



50 Celsius

- F. Equilibrium and Concentration
 - If the mass and temperature of a system remain constant, but the volume of the system increases, equilibrium will shift in order to maintain the concentrations of vapor particles
- II. Equilibrium Vapor Pressure of a Liquid
 - A. Equilibrium Vapor Pressure
 - 1. The pressure exerted by a vapor in equilibrium with its corresponding liquid at a given temperature
 - B. Volatile Liquids
 - 1. Liquids that have weak forces of attraction and evaporate easily
 - C. Nonvolatile Liquids
 - 1. Liquids that have strong forces of attraction and do not evaporate easily

III. Boiling

- A. Boiling
 - 1. The conversion of a liquid to a vapor within the liquid as well as at its surface. It occurs when the equilibrium vapor pressure of the liquid equals the atmospheric pressure
- B. Boiling Point
 - 1. The temperature at which the equilibrium vapor pressure of the liquid equals the atmospheric pressure
 - a. Water boils at 100 °C at 1 atm pressure
 - b. Water boils above 100 °C at higher pressures
 - c. Water boils below 100 °C at lower pressures
- C. Molar Heat of Vaporization
 - 1. The amount of heat energy required to vaporize one mole of a liquid at its boiling point
 - 2. Strong attractive forces between particles result in high molar heat of vaporization

IV. Freezing and Melting

- A. Freezing Point
 - 1. The temperature at which the solid and liquid are in equilibrium at 1 atm
 - 2. For pure crystalline solids, the melting point and freezing point are the same
 - 3. Temperature remains constant during a phase change
- B. Molar Heat of Fusion
 - 1. The amount of heat energy required to melt one mole of solid at its melting point
- C. Sublimation and Deposition
 - Sublimation is the change of state from a solid directly to a gas

 a. Dry ice → Gaseous CO₂
 - 2. Deposition is the change of state from a gas directly to a solid

- V. <u>Phase Diagrams</u>
 - A. Phase Diagram
 - 1. A graph of pressure versus temperature that shows the conditions under which the phases of a substance exist (notice that pressure is on a logarithmic scale)



- B. Triple Point
 - 1. The temperature and pressure conditions at which the solid, liquid, and vapor of the substance can coexist at equilibrium

C. Critical Temperature

- 1. The temperature at above which the substance cannot exist in the liquid state, regardless of pressure
 - a. For water, the critical temperature is 373.99 °C
- D. Critical Pressure
 - 1. The lowest pressure at which the substance can exist as a liquid at the critical temperature
 - a. For water, the critical pressure is 217.75 atm
- E. Critical Point
 - 1. The point on the graph describing simultaneously the critical temperature and the critical pressure

P = 217.75 atm

Temperature = 373.99 °C

12-4 Water

Bond Type	Polar	Density of ice (0 °C)	0.917 g/cm ³
Bond angle	105°	Density of water (0 °C)	0.999 g/cm ³
Boiling point	100 °	Point of maximum density	3.98 °C
Melting Point	0 °C	Molar heat of fusion	6.009 kJ/mole
		Molar heat of vaporization	40.79 kJ/mole